

Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Original) A method for preserving antigen presentation on a virally infected mammalian cell, comprising:

(a) providing a population of mammalian cells at least a portion of which is suspected of being virally infected and

(b) contacting said cells with an anti-apoptotic reagent, thereby preserving antigen presentation on virally infected cells.

2. (Original) The method of claim 1, wherein said cells comprise peripheral blood leukocytes.


3. (Original) The method of claim 1, wherein said cells comprise neutrophils.

4. (Original) The method of claim 1, wherein said cells comprise granulocytes.

5. (Original) The method of claim 1, wherein said virus is selected from the group consisting of herpes, HIV, cytomegalovirus (CMV), and hepatitis.

6. (Original) The method of claim 5, wherein said virus is CMV.

7. (Original) The method of claim 1, wherein said antigen comprises a viral antigen present on the surface of said mammalian cells.
8. (Original) The method of claim 7, wherein said antigen comprises pp65 protein of CMV.
9. (Original) The method of claim 1, wherein the contacting is *ex vivo*.
10. – 11. (Canceled)
12. (Original) The method of claim 1, wherein the reagent is a protease inhibitor.
13. (Original) The method of claim 12, wherein the protease inhibitor is irreversible.
14. (Original) The method of claim 12, wherein the protease inhibitor is reversible.
15. (Original) The method of claim 12, wherein the protease inhibitor is a compound of formula 1:



FORMULA 1

wherein:

n is 1 or 2;

R¹ is alkyl, cycloalkyl, (cycloalkyl)alkyl, phenyl, (substituted)phenyl, phenylalkyl, (substituted)phenylalkyl, heteroaryl, (heteroaryl)alkyl or (CH₂)_mCO₂R⁴, wherein m = 1-4, and R⁴ is as defined below;

R² is a hydrogen atom, chloro, alkyl, cycloalkyl, (cycloalkyl)alkyl, phenyl, (substituted)phenyl, phenylalkyl, (substituted)phenylalkyl, heteroaryl, (heteroaryl)alkyl or (CH₂)_pCO₂R⁵, wherein p = 0-4, and R⁵ is as defined below;

R³ is a hydrogen atom, alkyl, cycloalkyl, (cycloalkyl)alkyl, phenylalkyl, or (substituted)phenylalkyl;

R⁴ is a hydrogen atom, alkyl, cycloalkyl, (cycloalkyl)alkyl, phenylalkyl, or (substituted)phenylalkyl;

R⁵ is a hydrogen atom, alkyl, cycloalkyl, (cycloalkyl)alkyl, phenylalkyl, or (substituted)phenylalkyl;

A is a natural and unnatural amino acid;

B is a hydrogen atom, a deuterium atom, alkyl, cycloalkyl, (cycloalkyl)alkyl, phenyl, (substituted)phenyl, phenylalkyl, (substituted)phenylalkyl, heteroaryl, (heteroaryl)alkyl, halomethyl, CH₂ZR⁶, CH₂OCO(aryl), CH₂OCO(heteroaryl); or CH₂OPO(R⁷)R⁸, where Z is an oxygen or a sulfur atom;

R⁶ is phenyl, substituted phenyl, phenylalkyl, substituted phenylalkyl, heteroaryl, or (heteroaryl)alkyl; and


R⁷ and R⁸ are independently selected from a group consisting of alkyl, cycloalkyl, phenyl, substituted phenyl, phenylalkyl, (substituted phenyl) alkyl, and (cycloalkyl) alkyl; and

X and Y are independently selected from the group consisting of a hydrogen atom, halo, trihalomethyl, amino, protected amino, an amino salt, mono-substituted amino, di-substituted amino, carboxy, protected carboxy, a carboxylate salt, hydroxy, protected hydroxy, a salt of a hydroxy group, lower alkoxy, lower alkylthio, alkyl, substituted alkyl,

cycloalkyl, substituted cycloalkyl, (cycloalkyl)alkyl, substituted (cycloalkyl)alkyl, phenyl, substituted phenyl, phenylalkyl, and (substituted phenyl)alkyl;

or a pharmaceutically acceptable salt thereof.

16. (Original) The method of claim 12, wherein the protease inhibitor is a compound of formula 3:



FORMULA 3


wherein:

n is 1 or 2;

m is 1 or 2;

A is R^2CO- , $R^3-O-CO-$, or R^4SO_2- ;

a group of the formula:



further wherein:

R^1 is a hydrogen atom, alkyl or phenylalkyl;

R^2 is alkyl, cycloalkyl, (cycloalkyl)alkyl, phenyl, phenylalkyl, substituted phenyl, (substituted phenyl)alkyl, heteroaryl, or (heteroaryl)alkyl;

R^3 is alkyl, cycloalkyl, (cycloalkyl)alkyl, phenylalkyl, or (substituted phenyl)alkyl;


R^4 is alkyl, cycloalkyl, (cycloalkyl)alkyl, phenyl, phenylalkyl, substituted phenyl, (substituted phenyl)alkyl, heteroaryl, or (heteroaryl)alkyl;

R^5 is alkyl, cycloalkyl, (cycloalkyl)alkyl, phenyl, phenylalkyl, substituted phenyl, (substituted phenyl)alkyl, heteroaryl, or (heteroaryl)alkyl;

R^6 is alkyl, cycloalkyl, (cycloalkyl)alkyl, phenylalkyl, or (substituted phenyl)alkyl;

R^7 is alkyl, cycloalkyl, (cycloalkyl)alkyl, phenyl, phenylalkyl, substituted phenyl, (substituted phenyl)alkyl, heteroaryl, or (heteroaryl)alkyl;

R^8 is an amino acid side chain chosen from the group consisting of natural and unnatural amino acids;


 B is a hydrogen atom, a deuterium atom, alkyl, cycloalkyl, (cycloalkyl)alkyl, phenyl, phenylalkyl, substituted phenyl, (substituted phenyl)alkyl, heteroaryl, (heteroaryl)alkyl, or halomethyl;

a group of the formula:




wherein R^9 is phenyl, substituted phenyl, phenylalkyl, (substituted phenyl)alkyl, heteroaryl, or (heteroaryl)alkyl; and X is an oxygen or a sulfur atom;


a group of the formula:



a group of the formula:




a group of the formula:



wherein R^{10} and R^{11} are independently selected from a group consisting of alkyl, cycloalkyl, phenyl, substituted phenyl, phenylalkyl and (substituted phenyl) alkyl; and the pharmaceutically-acceptable salts thereof.

17. (Original) The method of claim 12, wherein the protease inhibitor is a compound of the formula:




Formula I

wherein:


A is a natural or unnatural amino acid of Formula IIa-i:




IIa




IIb




IIc




IIId




IIe




IIIf



IIg




IIh




IIi


B is a hydrogen atom, a deuterium atom, C₁₋₁₀ straight chain or branched alkyl, cycloalkyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, 2-benzoxazolyl, substituted 2-oxazolyl, (CH₂)_ncycloalkyl, (CH₂)_nphenyl, (CH₂)_n(substituted phenyl), (CH₂)_n(1 or 2-naphthyl), (CH₂)_n(heteroaryl), halomethyl, CO₂R¹², CONR¹³R¹⁴, CH₂ZR¹⁵, CH₂OCO(aryl), CH₂OCO(heteroaryl), or CH₂OPO(R¹⁶)R¹⁷, where Z is an oxygen or a sulfur atom, or B is a group of the Formula IIIa-c:



IIIa



IIIb



IIIc

A *1*

R^1 is alkyl, cycloalkyl, (cycloalkyl)alkyl, phenyl, substituted phenyl, phenylalkyl, substituted phenylalkyl, naphthyl, substituted naphthyl, (1 or 2 naphthyl)alkyl, heteroaryl, (heteroaryl)alkyl, $R^{1a}(R^{1b})N$, [or] $R^{1c}O$, 2-phenoxyphenyl or 2- or 3- benzylphenyl; and

R^2 is hydrogen, lower alkyl, cycloalkyl, (cycloalkyl)alkyl, phenylalkyl, or substituted phenylalkyl;

and wherein:

R^{1a} and R^{1b} are independently hydrogen, alkyl, cycloalkyl, (cycloalkyl)alkyl, phenyl, substituted phenyl, phenylalkyl, substituted phenylalkyl, naphthyl, substituted naphthyl, (1 or 2 naphthyl)alkyl, heteroaryl, or (heteroaryl)alkyl, with the proviso that R^{1a} and R^{1b} cannot both be hydrogen;

R^{1c} is alkyl, cycloalkyl, (cycloalkyl)alkyl, phenyl, substituted phenyl, phenylalkyl, substituted phenylalkyl, naphthyl, substituted naphthyl, (1 or 2 naphthyl)alkyl, heteroaryl, or (heteroaryl)alkyl;

R^3 is C_{1-6} lower alkyl, cycloalkyl, phenyl, substituted phenyl, $(CH_2)_nNH_2$, $(CH_2)_nNHCOR^9$, $(CH_2)_nN(C=NH)NH_2$, $(CH_2)_mCO_2R^2$, $(CH_2)_mOR^{10}$, $(CH_2)_mSR^{11}$, $(CH_2)_ncycloalkyl$, $(CH_2)_nphenyl$, $(CH_2)_n(substituted\ phenyl)$, $(CH_2)_n(1\ or\ 2-naphthyl)$ or $(CH_2)_n(heteroaryl)$, wherein heteroaryl includes pyridyl, thienyl, furyl, thiazolyl, imidazolyl,

pyrazolyl, isoxazolyl, pyrazinyl, pyrimidyl, triazinyl, tetrazolyl, and indolyl;

R^{3a} is hydrogen or methyl, or R^3 and R^{3a} taken together are $-(CH_2)_d-$ where d is an interger from 2 to 6;

R^4 is phenyl, substituted phenyl, $(CH_2)_m$ phenyl, $(CH_2)_m$ (substituted phenyl), cycloalkyl, or benzofused cycloalkyl;

R^5 is hydrogen, lower alkyl, cycloalkyl, phenyl, substituted phenyl, $(CH_2)_n$ cycloalkyl, $(CH_2)_n$ phenyl, $(CH_2)_n$ (substituted phenyl), or $(CH_2)_n$ (1 or 2-naphthyl);

R^6 is hydrogen, fluorine, oxo, lower alkyl, cycloalkyl, phenyl, substituted phenyl, naphthyl, $(CH_2)_n$ cycloalkyl, $(CH_2)_n$ phenyl, $(CH_2)_n$ (substituted phenyl), $(CH_2)_n$ (1 or 2-naphthyl), OR^{10} , SR^{11} or $NHCOR^9$;

R^7 is hydrogen, oxo, lower alkyl, cycloalkyl, phenyl, substituted phenyl, naphthyl, $(CH_2)_n$ cycloalkyl, $(CH_2)_n$ phenyl, $(CH_2)_n$ (substituted phenyl), or $(CH_2)_n$ (1 or 2-naphthyl);

R^8 is lower alkyl, cycloalkyl, $(CH_2)_n$ cycloalkyl, $(CH_2)_n$ phenyl, $(CH_2)_n$ (substituted phenyl), $(CH_2)_n$ (1 or 2-naphthyl), or COR^9 ;

R^9 is hydrogen, lower alkyl, cycloalkyl, phenyl, substituted phenyl, naphthyl, $(CH_2)_n$ cycloalkyl, $(CH_2)_n$ phenyl, $(CH_2)_n$ (substituted phenyl), $(CH_2)_n$ (1 or 2-naphthyl), OR^{12} , or $NR^{13}R^{14}$;

R^{10} is hydrogen, lower alkyl, cycloalkyl, phenyl, substituted phenyl, naphthyl, $(CH_2)_n$ cycloalkyl, $(CH_2)_n$ phenyl, $(CH_2)_n$ (substituted phenyl), or $(CH_2)_n$ (1 or 2-naphthyl);

R^{11} is lower alkyl, cycloalkyl, phenyl, substituted phenyl, naphthyl, $(CH_2)_n$ cycloalkyl, $(CH_2)_n$ phenyl, $(CH_2)_n$ (substituted phenyl), or $(CH_2)_n$ (1 or 2-naphthyl);

R^{12} is lower alkyl, cycloalkyl, $(CH_2)_n$ cycloalkyl, $(CH_2)_n$ phenyl, $(CH_2)_n$ (substituted phenyl), or $(CH_2)_n$ (1 or 2-naphthyl);

R^{13} is hydrogen, lower alkyl, cycloalkyl, phenyl, substituted phenyl, naphthyl, substituted naphthyl, $(CH_2)_n$ cycloalkyl, $(CH_2)_n$ phenyl, $(CH_2)_n$ (substituted phenyl), or $(CH_2)_n$ (1 or 2-naphthyl);

R^{14} is hydrogen or lower alkyl;

or R^{13} and R^{14} taken together form a five to seven membered carbocyclic or heterocyclic ring, such as morpholine, or N-substituted piperazine;

R^{15} is phenyl, substituted phenyl, naphthyl, substituted naphthyl, heteroaryl, $(CH_2)_n$ phenyl, $(CH_2)_n$ (substituted phenyl), $(CH_2)_n$ (1 or 2-naphthyl), or $(CH_2)_n$ (heteroaryl);

R^{16} and R^{17} are independently lower alkyl, cycloalkyl, phenyl, substituted phenyl, naphthyl, phenylalkyl, substituted phenylalkyl, or (cycloalkyl)alkyl;

R^{18} and R^{19} are independently hydrogen, alkyl, phenyl, substituted phenyl, $(CH_2)_n$ phenyl, $(CH_2)_n$ (substituted phenyl), or R^{18} and R^{19} taken together are $-(CH=CH)_2-$;

R^{20} is hydrogen, alkyl, phenyl, substituted phenyl, $(CH_2)_n$ phenyl, $(CH_2)_n$ (substituted phenyl);

R^{21} , R^{22} and R^{23} are independently hydrogen, or alkyl;

X is CH₂, (CH₂)₂, (CH₂)₃, or S;

Y¹ is O or NR²³;

Y² is CH₂, O, or NR²³;

a is 0 or 1 and b is 1 or 2, provided that when a is 1 then b is 1;

c is 1 or 2, provided that when c is 1 then a is 0 and b is 1;

m is 1 or 2; and

n is 1, 2, 3 or 4;

or a pharmaceutically acceptable salt thereof.

Al
cond'd